

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is designed for cleaning, deodorizing, and disinfecting for hospitals, nursing homes, schools, hotels, homes, and food processing establishments. This product is effective where housekeeping is of prime importance in controlling cross-contamination from treated surfaces.

This product when used as directed is formulated to disinfect inanimate hard non-porous surfaces: walls, floors, sink tops, toilet bowls, toilet seats, chairs, tables, telephones, restrooms, shower stalls, bathtubs, hampers, and cabinets.

This product is not for use on medical device surfaces.

This product deodorizes those areas which generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, toilet bowls, and other areas which are prone to odors caused by microorganisms.

DISINFECTION, DEODORIZATION AND CLEANING

- Shake can before using. Hold container six to eight inches from surface to be treated. Spray area until it is covered with white foam. Allow foam to penetrate and remain wet for 10 minutes. No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. This product will not leave grit or soap scum.

For heavily soiled areas, a pre-cleaning step is required before treatment.

When used in food processing establishments, cover or remove all food and packaging materials, remove all gross soils, saturate all surfaces with the spray product, scrub to loosen all soils, allow to soak for 10 minutes, thoroughly rinse all wetted and cleaned surfaces with potable water.

MILDEWSTAT - To control mold and mildew on pre-cleaned, hard, non-porous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth appears.

FUNGICIDE - This product is fungicidal against the pathogenic fungi, Trichophyton mentagrophytes (Athlete's Foot Fungus), when used as directed on hard surfaces found in bathrooms, shower stalls, locker rooms, or other clean, non-porous, hard surfaces commonly contacted by bare feet.

TUBERCULOCIDAL ACTIVITY - This product exhibits disinfectant efficacy against Mycobacterium tuberculosis (BCG) in 10 minutes at 20 degrees Centigrade when used as directed on previously cleaned hard nonporous inanimate surfaces.

BACTERICIDAL ACTIVITY - When used as directed, this product exhibits effective disinfectant activity against the organisms: Staphylococcus aureus, Salmonella choleraesuis, Pseudomonas aeruginosa, Escherichia coli 0157:H7, Methicillin resistant Staphylococcus aureus (MRSA), Vancomycin intermediate resistant Staphylococcus aureus (VISA), Vancomycin resistant Enterococcus faecalis (VRE) and meets the requirements for hospital use.

***VIRUCIDAL ACTIVITY** - This product, when used on environmental, inanimate, non-porous surfaces, exhibits effective virucidal activity against HIV-1 (associated with AIDS), Canine Parvovirus, Poliovirus Type 1, Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV).



Lift Off™

FOAMING GERMICIDAL CLEANER

ACTIVE INGREDIENTS

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium chlorides.....	0.105%
n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl ethylbenzyl ammonium chlorides.....	0.105%
INERT INGREDIENTS.....	99.790%
TOTAL.....	100.000%

KEEP OUT OF REACH OF CHILDREN WARNING

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator. See side panel for additional precautionary statements.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING. KEEP OUT OF REACH OF CHILDREN. Causes eye and skin irritation. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

EPA REG. NO. 3862-181-34370
EPA EST. NO. 9143-TX-1

Distributed By:

WECHEM, INC.

5734 Susitna Dr., Harahan, LA 70123

(504) 733-1152 • 1-800-426-0512

FAX: (504) 733-2218

NET WT.: 19 OZ (1LB 3OZ)(539 G)

12/22

KILLS HIV-1, HBV, AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

PERSONAL PROTECTION: When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks, or eye coverings.

CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

CONTACT TIME: Allow surface to remain wet for 10 minutes.

DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE, FUNGICIDE, AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

CONTAINER DISPOSAL - Replace cap and dispose of container in trash. Do not incinerate or puncture. **STORAGE** - Store in a dry place no lower in temperature than 50° F or higher than 120° F.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. 472420F5788.122822



SCAN FOR SDS